arXiv:1412.0764v1 [math.DS] 2 Dec 2014

Large Deviation Properties in Some Nonuniformly Hyperbolic Systems Via Pesin Theory

Zheng Yin[1], Ercai Chen*[1,2]

1 School of Mathematical Sciences and Institute of Mathematics, Nanjing Normal University,

Nanjing 210023, Jiangsu, P.R.China

2 Center of Nonlinear Science, Nanjing University,

Nanjing 210093, Jiangsu, P.R.China.

e-mail: zhengyinmail@126.com ecchen@njnu.edu.cn

Abstract. This article is devoted to level-1 large deviation properties in some nonuniformly hyperbolic systems via Pesin theory. In particular, our result can be applied to the nonuniformly hyperbolic diffeomorphisms described by Katok and several other classes of diffeomorphisms derived from Anosov systems.

Keywords and phrases: Large deviations, Pesin set, weak shadowing property.

1 Introduction

The classical theory of large deviations deals with the rates of convergence of sequence of random variables to some limit distribution. We refer to Ellis' book [8] for a general theory of large deviations and its background in statistical mechanics. We consider large deviations for dynamical systems. Let (M, d, f) be a topological dynamical system and let μ be a ergodic measure. Given an observable $\varphi \in L^1(\mu)$, let $S_n\varphi(x) = \sum_{i=0}^{n-1} \varphi(f^i x)$. By Birkhoff ergodic theorem, $(1/n)S_n\varphi(x) \to \int \varphi d\mu$ as $n \to \infty$ for almost all $x \in M$. We are interested in the rate of convergence of this time average. More generally, given a reference measure m, one interesting question is to study the asymptotic behaviour of

$$m\left\{x \in M : \frac{1}{n}S_n\varphi(x) \in E\right\}$$

* Corresponding author

2010 Mathematics Subject Classification: 37D25, 37C40, 60F10

for some $E \subseteq \mathbb{R}$. This topic had contributions by many authors in the recent years. We refer the reader to [1, 5, 6, 7, 13, 15, 18, 20, 21, 22, 24, 25] and the references therein.

The large deviation principle can be divided into two steps, i.e., to obtain the upper bound of

$$\limsup_{n\to\infty} \frac{1}{n} \log m \left\{ x \in M : \frac{1}{n} S_n \varphi(x) \in E \right\}$$

and the lower bound of

$$\liminf_{n\to\infty} \frac{1}{n} \log m \left\{ x \in M : \frac{1}{n} S_n \varphi(x) \in E \right\}. \tag{1.1}$$

The upper bound is usually easier to establish than the lower bound. An effective method in dynamical systems to obtain the lower estimate for (1.1) is "orbit-gluing approach". For example, Young [24] formulated quite general large deviation for dynamical systems via specification property. Pfister and Sullivan [18] introduced approximate product property to study the large deviations estimates for β-shifts. Yamamoto [23] considered the large deviation principle for group automorphisms which satisfy the almost specification property. Varandas [22] introduced a measure theoretical nonuniform specification to obtain large deviations estimstes for weak Gibbs measures.

Recently, many efforts have been made to extend the theory of large deviations to the scope of nonuniformly hyperbolic systmes. Araújo and Pacífico [1] gave lower as well as upper bounds in the noninvertible context. Melbourne and Nicol [15] gave optimal bounds in the Young tower situation (both invertible and noninvertible). Large dviations principles were also obtained by Yuri [25] in the context shifts with countably many symbols and by Comman and Rivera [6] for nonuniformly expanding rational maps. More recently, Chung [5] obtained a sufficient condition to hold a full large deviation principle for Markov tower maps induced from return time functions and also gave counterexamples when [6]'s conditions do not hold.

Motivated by the work of Young [24] and Varandas [22] our purpose here is to give a contribution for large deviations estimates in some nonuniformly hyperbolic systems. The main tool we use is weak shadowing lemma in Pesin theory. Particularly, our result can be applied (i) to the diffeomorphisms on surfaces, (ii) to the nonuniformly hyperbolic diffeomorphisms described by Katok and several other classes of diffeomorphisms derived from Anosov systems.

This article is organized as follows. In section 2, we provide some notions and results of Pesin theory and state the main result. Section 3 is devoted to the proof of the main result. Examples and applications are given in section 4.

2 Preliminaries

In this section, we first present some notations to be used in this paper. Then we introduce some notions and results of Pesin theory [3, 12, 19] and state the main results.

We denote by $\mathscr{M}(M,f)$, $\mathscr{M}_{\rm inv}(M,f)$ and $\mathscr{M}_{\rm erg}(M,f)$ the set of all Borel probability measures, f-invariant Borel probability measures and ergodic measures respectively. It is well known that $\mathscr{M}(M,f)$ equiped with weak* topology is a compact metrisable space. We will denote by D the metric on $\mathscr{M}(M,f)$. For an f-invariant subset $Z\subset X$, let $\mathscr{M}_{\rm inv}(Z,f)$ denote the subset of $\mathscr{M}_{\rm inv}(M,f)$ for which the measures μ satisfy $\mu(Z)=1$ and $\mathscr{M}_{\rm erg}(Z,f)$ denote those which are ergodic. Denote by $C^0(M)$ the space of continuous functions from M to $\mathbb{R}$ with the sup norm. For $\varphi\in C^0(M)$ and $n\geq 1$ we denote $\sum_{i=0}^{n-1}\varphi(f^ix)$ by $S_n\varphi(x)$. For every $\epsilon>0$, $n\in\mathbb{N}$ and a point $x\in M$, define $B_n(x,\epsilon)=\{y\in M: d(f^ix,f^iy)<\epsilon, \forall 0\leq i\leq n-1\}$.

Suppose M is a compact connected boundary-less Riemannian n-dimension manifold and $f:X\to X$ is a $C^{1+\alpha}$ diffeomorphism. Let $\mu\in\mathscr{M}_{\rm erg}(Z,f)$ and Df_x denote the tangent map of f at $x\in M$. We say that $x\in X$ is a regular point of f if there exist $\lambda_1(\mu)>\lambda_2(\mu)>\cdots>\lambda_{\phi(\mu)}(\mu)$ and a decomposition on the tangent space $T_xM=E_1(x)\oplus\cdots\oplus E_{\phi(\mu)}(x)$ such that

$$\lim_{n\to\infty}\frac{1}{n}\log\|(Df_x^n)u\|=\lambda_j(x),$$

where $0\neq u\in E_j(x), 1\leq j\leq\phi(\mu)$. The number $\lambda_j(x)$ and the space $E_j(x)$ are called the Lyapunov exponents and the eigenspaces of f at the regular point x, respectively. Oseledets theorem [16] say that all regular points forms a Borel set with total measure. For a regular point $x\in M$, we define

$$\lambda^+(\mu)=\min\{\lambda_i(\mu)|\lambda_i(\mu)\geq 0, 1\leq i\leq\phi(\mu)\}$$

and

$$\lambda^-(\mu)=\min\{-\lambda_i(\mu)|\lambda_i(\mu)\leq 0, 1\leq i\leq\phi(\mu)\}.$$

We appoint $\min\emptyset=0$. An ergodic measure ω is hyperbolic if $\lambda^+(\omega)$ and $\lambda^-(\omega)$ are both non-zero.

Definition 2.1. *Given $\beta_1,\beta_2\gg\epsilon>0$ and for all $k\in\mathbb{Z}^+$, the hyperbolic block $\Lambda_k=\Lambda_k(\beta_1,\beta_2,\epsilon)$ consists of all points $x\in M$ such that there exists a decomposition $T_xM=E_x^s\oplus E_x^u$ satisfying:*

- $Df^t(E_x^s)=E_{f^tx}^s$ *and* $Df^t(E_x^u)=E_{f^tx}^u$;
- $\|Df^n|E_{f^tx}^s\|\leq e^{\epsilon k}e^{-(\beta_1-\epsilon)n}e^{\epsilon|t|}, \forall t\in\mathbb{Z}, n\geq 1$;

- $\|Df^{-n}|E^u_{f^tx}\| \le e^{\epsilon k}e^{-(\beta_2-\epsilon)n}e^{\epsilon|t|}, \forall t \in \mathbb{Z}, n \ge 1$;
- $\tan(\angle(E^s_{f^tx}, E^u_{f^tx})) \ge e^{-\epsilon k}e^{-\epsilon|t|}, \forall t \in \mathbb{Z}$.

Definition 2.2. $\Lambda = \Lambda(\beta_1, \beta_2, \epsilon) = \bigcup_{k=1}^{\infty} \Lambda_k(\beta_1, \beta_2, \epsilon)$ *is a Pesin set.*

The following statements are elementary properties of Pesin blocks (see [19]):

(1) $\Lambda_1 \subseteq \Lambda_2 \subseteq \cdots$;

(2) $f(\Lambda_k) \subseteq \Lambda_{k+1}, f^{-1}(\Lambda_k) \subseteq \Lambda_{k+1}$;

(3) Λ_k is compact for each $k \ge 1$;

(4) For each $k \ge 1$, the splitting $\Lambda_k \ni x \mapsto E^s_x \oplus E^u_x$ is continuous.

The Pesin set $\Lambda(\beta_1, \beta_2, \epsilon)$ is an f-invariant set but usually not compact. Given an ergodic measure $\mu \in \mathscr{M}_{\text{erg}}(M, f)$, denote by $\mu|\Lambda_l$ the conditional measure of μ on Λ_l. Let $\widetilde{\Lambda}_l = \text{supp}(\mu|\Lambda_l)$ and $\widetilde{\Lambda}_\mu = \bigcup_{l\ge1} \widetilde{\Lambda}_l$. If ω is an ergodic hyperbolic measure for f and $\beta_1 \le \lambda^-(\omega)$ and $\beta_2 \le \lambda^+(\omega)$, then $\omega \in \mathscr{M}_{\text{inv}}(\widetilde{\Lambda}_\omega, f)$.

Let $\{\delta_k\}_{k=1}^{\infty}$ be a sequence of positive real numbers. Let $\{x_n\}_{n=-\infty}^{\infty}$ be a sequence of points in $\Lambda = \Lambda(\beta_1, \beta_2, \epsilon)$ for which there exists a sequence $\{s_n\}_{n=-\infty}^{\infty}$ of positive integers satisfying:

$$\begin{aligned}&\text{(a) } x_n \in \Lambda_{s_n}, \forall n \in \mathbb{Z};\\ &\text{(b) } |s_n - s_{n-1}| \le 1, \forall n \in \mathbb{Z};\\ &\text{(c) } d(f(x_n), x_{n+1}) \le \delta_{s_n}, \forall n \in \mathbb{Z},\end{aligned}$$

then we call $\{x_n\}_{n=-\infty}^{\infty}$ a $\{\delta_k\}_{k=1}^{\infty}$ pseudo-orbit. Given $\eta > 0$ a point $x \in M$ is an η-shadowing point for the $\{\delta_k\}_{k=1}^{\infty}$ pseudo-orbit if $d(f^n(x), x_n) \le \eta\epsilon_{s_n}, \forall n \in \mathbb{Z}$, where $\epsilon_k = \epsilon_0 e^{-\epsilon k}$ and ϵ_0 is a constant only dependent on the system of f.

Weak shadowing lemma. [9, 12, 19] *Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism, with a non-empty Pesin set $\Lambda = \Lambda(\beta_1, \beta_2, \epsilon)$ and fixed parameters, $\beta_1, \beta_2 \gg \epsilon > 0$. For $\eta > 0$ there exists a sequence $\{\delta_k\}$ such that for any $\{\delta_k\}$ pseudo orbit there exists a unique η-shadowing point.*

Let m be a finite Borel measure on M. We think of m as our reference measure. We define

$$\mathcal{V}^+ = \big\{\psi \in C^0(M) : \exists C, \epsilon > 0 \text{ such that } \forall x \in M \text{ and } \forall n \ge 0, \\ mB_n(x, \epsilon) \le Ce^{-S_n\psi(x)}\big\}$$

and

$$\mathcal{V}^- = \big\{\psi \in C^0(M) : \exists \text{ arbitrarily small } \epsilon > 0 \text{ and } C = C(\epsilon) > 0 \\ \text{such that } \forall x \in M \text{ and } \forall n \geq 0, mB_n(x,\epsilon) \geq Ce^{-S_n\psi(x)}\big\}.$$

For $\varphi \in C^0(M)$ and $E \subseteq \mathbb{R}$, we write

$$\overline{R}(\varphi, E) = \limsup_{n\to\infty} \frac{1}{n} \log m \left\{ x \in M : \frac{1}{n} S_n\varphi(x) \in E \right\}$$

and

$$\underline{R}(\varphi, E) = \liminf_{n\to\infty} \frac{1}{n} \log m \left\{ x \in M : \frac{1}{n} S_n\varphi(x) \in E \right\}.$$

Now, we state the main result of this paper as follows:

Theorem 2.1. *Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism of a compact Riemannian manifold, with a non-empty Pesin set $\Lambda = \Lambda(\beta_1, \beta_2, \epsilon)$ and fixed parameters, $\beta_1, \beta_2 \gg \epsilon > 0$. Then for every $\varphi \in C^0(M)$ and $c \in \mathbb{R}$, the following hold:*

(1) For $\psi \in \mathcal{V}^+$, we have

$$\overline{R}(\varphi, [c, \infty)) \leq \sup \left\{ h_\nu(f) - \int \psi d\nu : \nu \in \mathscr{M}_{\text{inv}}(M, f), \int \varphi d\nu \geq c \right\}.$$

(2) For $\psi \in \mathcal{V}^-$ and $\mu \in \mathscr{M}_{erg}(M, f)$, we have

$$\underline{R}(\varphi, (c, \infty)) \geq \sup \left\{ h_\nu(f) - \int \psi d\nu : \nu \in \mathscr{M}_{inv}(\widetilde{\Lambda}_\mu, f), \int \varphi d\nu > c \right\}.$$

Corollary 2.1. *Let $f : M \to M$ be a $C^{1+\alpha}$ diffeomorphism of a compact Riemannian manifold and let $\omega \in \mathscr{M}_{\text{erg}}(M, f)$ be a hyperbolic measure. If $\beta_1 \leq \lambda^-(\omega)$ and $\beta_2 \leq \lambda^+(\omega)$, then for every $\varphi \in C^0(M)$ and $c \in \mathbb{R}$, the following hold:*

(1) For $\psi \in \mathcal{V}^+$, we have

$$\overline{R}(\varphi, [c, \infty)) \leq \sup \left\{ h_\nu(f) - \int \psi d\nu : \nu \in \mathscr{M}_{\text{inv}}(M, f), \int \varphi d\nu \geq c \right\}.$$

(2) For $\psi \in \mathcal{V}^-$, we have

$$\underline{R}(\varphi, (c, \infty)) \geq \sup \left\{ h_\nu(f) - \int \psi d\nu : \nu \in \mathscr{M}_{inv}(\widetilde{\Lambda}_\omega, f), \int \varphi d\nu > c \right\},$$

where $\widetilde{\Lambda}_\omega = \bigcup_{l\geq 1} \text{supp}(\omega|\Lambda_l(\beta_1, \beta_2, \epsilon))$.

3 Proof of Main Theorem

In this section, we will verify theorem 2.1. The upper bound for the measure of deviation holds for all topological dynamical systems. By [24], for $\psi \in \mathcal{V}^+$ we have

$$\overline{R}(\varphi, [c, \infty)) \leq \sup \left\{ h_\nu(f) - \int \psi d\nu : \nu \in \mathscr{M}_{\text{inv}}(M, f), \int \varphi d\nu \geq c \right\}.$$

To obtain the lower bound estimate we need to construct a suitable pseudo-orbit. Our method is inspired by [14] and [24]. Firstly, we give some important lemmas as follows.

Lemma 3.1. [11] *Let (X, d) be a compact metric space, $f : X \to X$ be a continuous map and μ be an ergodic invariant measure. For $\epsilon > 0$, $\gamma \in (0, 1)$ let $N^\mu(n, \epsilon, \gamma)$ denote the minimum number of ϵ-Bowen balls $B_n(x, \epsilon)$, which cover a set of μ-measure at least $1 - \gamma$. Then*

$$h_\mu(f) = \lim_{\epsilon \to 0} \liminf_{n \to \infty} \frac{1}{n} \log N^\mu(n, \epsilon, \gamma) = \lim_{\epsilon \to 0} \limsup_{n \to \infty} \frac{1}{n} \log N^\mu(n, \epsilon, \gamma).$$

Lemma 3.2. [4] *Let $f : M \to M$ be a C^1 diffeomorphism of a compact Riemannian manifold and $\mu \in \mathscr{M}_{\text{inv}}(M, f)$. Let $\Gamma \subseteq M$ be a measurable set with $\mu(\Gamma) > 0$ and let*

$$\Omega = \bigcup_{n \in \mathbb{Z}} f^n(\Gamma).$$

Take $\gamma > 0$. Then there exists a measurable function $N_0 : \Omega \to \mathbb{N}$ such that for a.e.$x \in \Omega$ and every $t \in [0, 1]$ there is some $l \in \{0, 1, \cdots, n\}$ such that $f^l(x) \in \Gamma$ and $|(l/n) - t| < \gamma$.

Fix $\psi \in \mathcal{V}^-$. Pick an arbitrary $\nu \in \mathscr{M}_{inv}(\widetilde{\Lambda}, f)$ with $\int \varphi d\nu > c$. We will prove that

$$\liminf_{n \to \infty} \frac{1}{n} \log \left\{ x \in M : \frac{1}{n} S_n \varphi(x) \in E \right\} \geq h_\nu(f) - \int \psi d\nu - 10\gamma - \gamma h_{top}(f).$$

for any preassigned $0 < \gamma < 1$. Let $\delta = \frac{1}{5}(\int \varphi d\nu - c)$.

Lemma 3.3. *For $\gamma > 0, \delta > 0$ and $\nu \in \mathscr{M}_{inv}(\widetilde{\Lambda}, f)$ there exists a finite convex combination of ergodic probability measures with rational coefficients $\lambda = \sum_{i=1}^{k} a_i \mu_i$ such that*

$$\mu_i(\widetilde{\Lambda}) = 1, \int \varphi d\lambda \geq c + 4\delta, \int \psi d\nu \geq \int \psi d\lambda - \gamma \text{ and } h_\lambda(f) \geq h_\nu(f) - \gamma.$$

Proof. We choose $\epsilon_1 > 0$ sufficiently small such that for any $\mu_1, \mu_2 \in \mathscr{M}_{\text{inv}}(M, f)$ with $D(\mu_1, \mu_2) < \epsilon_1$ we have

$$\left| \int \varphi d\mu_1 - \int \varphi d\mu_2 \right| < \delta, \left| \int \psi d\mu_1 - \int \psi d\mu_2 \right| < \gamma.$$

By ergodic decomposition theorem, we can write $\nu = \int_{\mathscr{M}_{erg}(\widetilde{\Lambda},f)} m d\tau(m)$. Let $\mathscr{P} = \{P_1, P_2, \cdots, P_k\}$ be a partition of $\mathscr{M}_{erg}(\widetilde{\Lambda}, f)$ with $\mathrm{diam}\mathscr{P} < \epsilon_1$. Let $a_i = \tau(P_i)$. For every $1 \leq i \leq k$ we can choose an ergodic measure $\mu_i \in P_i$ satisfying $h_{\mu_i}(f) \geq h_m(f) - \gamma$ for τ-almost every $m \in P_i$. Let $\lambda = \sum_{i=1}^{k} a_i \mu_i$. Obviously $\mu_i(\widetilde{\Lambda}) = 1, 1 \leq i \leq k$. By ergodic decomposition theorem one can easily prove the remaining inqualities. Since the a_i's can be approximated by rational numbers, the desired results follows. □

Since $\mu_i(\widetilde{\Lambda}) = 1$, we can choose $l \in \mathbb{N}$ such that $\mu_i(\widetilde{\Lambda}_l) > 1 - \gamma$ for all $1 \leq i \leq k$. By lemma 3.1, we can choose ϵ' sufficiently small so

$$d(x, y) < \epsilon' \Rightarrow |\varphi(x) - \varphi(y)| < \delta, |\psi(x) - \psi(y)| < \gamma \tag{3.1}$$

and

$$\liminf_{n\to\infty} \frac{1}{n} \log N^{\mu_i}(n, 4\epsilon', \gamma) > h_{\mu_i}(f) - \gamma$$

for $1 \leq i \leq k$. Let $\eta = \frac{\epsilon'}{\epsilon_0}$, it follows from weak shadowing lemma that there is a sequence of numbers $\{\delta_k\}$. Let ξ be a finite partition of X with $\mathrm{diam}(\xi) < \frac{\delta_l}{3}$ and $\xi \geq \{\widetilde{\Lambda}_l, M \setminus \widetilde{\Lambda}_l\}$. For $1 \leq i \leq k, n \in \mathbb{N}$, we consider the set

$$\Lambda_1^n(\mu_i) = \left\{ x \in \widetilde{\Lambda}_l : f^q(x) \in \xi(x) \text{ for some } q \in [n, (1+\gamma)n] \right\}$$

and

$$\begin{aligned} \Lambda_2^n(\mu_i) = \Big\{ x \in \widetilde{\Lambda}_l : & S_m\psi(x) \leq m(\int \psi d\mu_i + \gamma) \text{ and} \\ & S_m\varphi(x) \geq m(\int \varphi d\mu_i - \delta) \text{ for all } m \geq n \Big\}, \end{aligned} \tag{3.2}$$

where $\xi(x)$ is the element in ξ containing x. Let $\Lambda^n(\mu_i) = \Lambda_1^n(\mu_i) \cap \Lambda_2^n(\mu_i)$. Then we have the following lemma.

Lemma 3.4. $\lim_{n\to\infty} \mu_i(\Lambda^n(\mu_i)) = \mu_i(\widetilde{\Lambda}_l), 1 \leq i \leq k.$

Proof. By Birkhoff ergodic theorem, it suffices to prove that

$$\lim_{n\to\infty} \mu_i(\Lambda_1^n(\mu_i)) = \mu_i(\widetilde{\Lambda}_l), 1 \leq i \leq k.$$

Let $[(1+\gamma)n]$ be the integer part of $(1+\gamma)n$. Take an element ξ_k of ξ and let $\Gamma = \xi_k$. Applying lemma 3.2, we can deduce that for $a.e.x \in \xi_k$, there exists a measurable function N_0 such that for every $[(1+\gamma)n] \geq N_0(x)$ there is some $q \in \{0, 1, \cdots, [(1+\gamma)n]\}$ such that $f^q(x) \in \xi_k$ and

$$\frac{q}{[(1+\gamma)n]} - 1 > -\frac{\gamma}{1+\gamma}.$$

We obtain $q \geq n - \frac{1}{1+\gamma}$. Since $q \in \mathbb{N}$, we have $q \in [n, (1+\gamma)n]$ and the proof is completed. □

By lemma 3.1, we can take sufficiently large $K \in \mathbb{N}$ such that

$$\mu_i(\Lambda^n(\mu_i)) > 1 - \gamma$$

for all $n \geq M$ and $1 \leq i \leq k$.

Let

$$Q(\Lambda^n(\mu_i), 4\epsilon') = \inf\{\sharp S : S \text{ is } (n, 4\epsilon') \text{ spanning set for } \Lambda^n(\mu_i)\},$$
$$P(\Lambda^n(\mu_i), 4\epsilon') = \sup\{\sharp S : S \text{ is } (n, 4\epsilon') \text{ separated set for } \Lambda^n(\mu_i)\}.$$

Then for all $n \geq K$ and $1 \leq i \leq k$, we have

$$P(\Lambda^n(\mu_i), 4\epsilon') \geq Q(\Lambda^n(\mu_i), 4\epsilon') \geq N^{\mu_i}(n, \epsilon', \gamma).$$

We obtain

$$\liminf_{n\to\infty} \frac{1}{n} \log P(\Lambda^n(\mu_i), 4\epsilon') \geq \liminf_{n\to\infty} \frac{1}{n} \log N^{\mu_i}(n, 4\epsilon', \gamma) > h_{\mu_i}(f) - \gamma.$$

Thus we can choose $t \in \mathbb{N}$ large enough such that $\exp(t\gamma) > \sharp\xi$ and

$$\frac{1}{t} \log P(\Lambda^t(\mu_i), 4\epsilon') > h_{\mu_i}(f) - 2\gamma.$$

for $1 \leq i \leq k$. Let S_i be a $(t, 4\epsilon')$-separated set for $\Lambda^t(\mu_i)$ and

$$\#S_i \geq \exp\left(t(h_{\mu_i}(f) - 3\gamma)\right).$$

For each $q \in [t, (1+\gamma)t]$, let

$$V_q(\mu_i) = \{x \in S_i : f^q(x) \in \xi(x)\}$$

and let n_i be the value of q which maximizes $\#V_q(\mu_i)$. Obviously, $n_i \geq t \geq \frac{n_i}{1+\gamma} \geq n_i(1-\gamma)$. Since $\exp(t\gamma) \geq t\gamma + 1$, we have that

$$\#V_{n_i}(\mu_i) \geq \frac{\#S_i}{t\gamma + 1} \geq \exp\left(t(h_{\mu_i}(f) - 4\gamma)\right).$$

Consider the element $A_i \in \xi$ such that $\#(V_{n_i}(\mu_i) \cap A_i)$ is maximal. Let $W_i = V_{n_i}(\mu_i) \cap A_i$. It follows that

$$\#W_i \geq \frac{1}{\#\xi} \#V_{n_i}(\mu_i) \geq \frac{1}{\#\xi} \exp\left(t(h_{\mu_i}(f) - 4\gamma)\right).$$

Since $e^{t\gamma} > \sharp\xi$, $t \geq n_i(1-\gamma)$ and $\#W_i \geq 1$, we have

$$\#W_i \geq \exp\left(n_i(1-\gamma)(h_{\mu_i}(f) - 5\gamma)\right).$$

Notice that A_i is contained in an open subset U_i with $\mathrm{diam}(U_i) \leq 3\mathrm{diam}(\xi)$. By the ergodicity of μ, for any two measures μ_{i_1}, μ_{i_2}, there exists $s = s(\mu_{i_1}, \mu_{i_2}) \in \mathbb{N}$ and $y = y(\mu_{i_1}, \mu_{i_2}) \in U_{i_1} \cap \widetilde{\Lambda}_l$ such that $f^s(y) \in U_{i_2} \cap \widetilde{\Lambda}_l$. Letting $C_i = \frac{a_i}{n_i}$, we can choose an integer N large enough so that NC_i, $1 \leq i \leq k$ are integers and

$$N\gamma \geq \sum_{i=1}^{k-1} s(\mu_i, \mu_{i+1}).$$

Let $X = \sum_{i=1}^{k-1} s(\mu_i, \mu_{i+1})$ and

$$Y_n = n \sum_{i=1}^{k} Nn_iC_i + X = nN + X, n = 1, 2, \cdots.$$

Then we have

$$\frac{X}{N} \leq \gamma, \frac{N}{Y_n} \geq \frac{1}{n+\gamma}, \frac{X}{Y_n} \leq \frac{\gamma}{n+\gamma}. \tag{3.3}$$

For simplicity of the statement, for $x \in M$ define segments of orbits

$$\begin{aligned} L_i(x) &:= (x, f(x), \cdots, f^{n_i-1}(x)), 1 \leq i \leq k, \\ \widehat{L}_{i_1,i_2}(x) &:= (x, f(x), \cdots, f^{s(\mu_{i_1},\mu_{i_2})-1}(x)), 1 \leq i_1, i_2 \leq k. \end{aligned}$$



Figure 1: Quasi-orbit

For every $n \in \mathbb{N}$, consider the pseudo-orbit with finite length (see Figure 1)

$$\begin{aligned} \Big\{ & L_1(x(1,1)), L_1(x(1,2)), \cdots, L_1(x(1, nNC_1)), \widehat{L}_{1,2}(y(\mu_1, \mu_2)); \\ & L_2(x(2,1)), L_2(x(2,2)), \cdots, L_2(x(2, nNC_2)), \widehat{L}_{2,3}(y(\mu_2, \mu_3)); \\ & \vdots \\ & L_k(x(k,1)), L_k(x(k,2)), \cdots, L_k(x(k, nNC_k)) \Big\}, \end{aligned}$$

where $x(i,j) \in W_i, 1 \leq i \leq k, 1 \leq j \leq nNC_i$.

For $1 \le i \le k$, $1 \le j \le nNC_i$, let

$$\begin{aligned} M_1 &= 0, \\ M_i &= \sum_{l=1}^{i-1} (nNC_l n_l + s(\mu_l, \mu_{l+1})), \\ M_{i,j} &= M_i + (j-1)n_i. \end{aligned}$$

By weak shadowing lemma, there exists a shadowing z for the above pseudo-orbit satisfying

$$d(f^{M_{i,j}+q}(z), f^q(x(i,j))) \le \eta\epsilon_0 e^{-\epsilon l} = \frac{\epsilon'}{\epsilon_0}\epsilon_0 e^{-\epsilon l} \le \epsilon'$$

for $0 \le q \le n_i - 1$. The shadowing point z can be considered as a map with the variables $x(i,j)$. We define F_n to be the image of the map. More precisely,

$$\begin{aligned} F_n = \Big\{ z : z = z(x(1,1), x(1,2), \cdots, x(1,nNC_1), x(2,1), x(2,2), \cdots, x(2,nNC_2), \cdots, \\ x(k,1), x(k,2), \cdots, x(k,nNC_k)), x(i,j) \in W_i, 1 \le i \le k, 1 \le j \le nNC_i \Big\}. \end{aligned}$$

Since W_i is $(n_i, 4\epsilon')$ separated sets, for distinct $x(i,j), x'(i,j) \in W_i$ the corresponding shadowing points $z, z' \in F_n$ satisfying $B_{Y_n}(z,\epsilon') \cap B_{Y_n}(z',\epsilon') = \emptyset$. Thus

$$\begin{aligned} \#F_n &\ge (\#W_1)^{nNC_1} (\#W_2)^{nNC_2} \cdots (\#W_k)^{nNC_k} \\ &\ge \exp\left\{ \sum_{i=1}^{k} nNn_i C_i (1-\gamma)(h_{\mu_i}(f) - 5\gamma) \right\} \\ &= \exp\left\{ nN(1-\gamma)(h_\lambda(f) - 5\gamma) \right\}. \end{aligned} \tag{3.4}$$

Lemma 3.5. *For sufficiently large n, we have $B_{Y_n}(z,\epsilon') \subseteq \left\{ x \in M : \frac{1}{Y_n} S_{Y_n}\varphi(x) > c \right\}$ for all $z \in F_n$.*

Proof. From (3.1), it suffices to prove that for sufficiently large n,

$$\frac{1}{Y_n} S_{Y_n}\varphi(z) > c + \delta.$$

By (3.1), (3.2), $a_i = C_i n_i$ and lemma 3.3, we have

$$\begin{aligned} \frac{1}{Y_n} S_{Y_n}\varphi(z) &\ge \frac{1}{Y_n} \sum_{i=1}^{k} \sum_{j=1}^{nNC_i} S_{n_i}\varphi(x(i,j)) - \frac{nN\delta}{Y_n} - \frac{X\|\varphi\|}{Y_n} \\ &\ge \frac{1}{Y_n} \sum_{i=1}^{k} nNC_i n_i \Big(\int \varphi d\mu_i - \delta \Big) - \frac{nN\delta}{Y_n} - \frac{X\|\varphi\|}{Y_n} \\ &= \frac{nN}{Y_n} \Big(\int \varphi d\lambda - \delta \Big) - \frac{nN\delta}{Y_n} - \frac{X\|\varphi\|}{Y_n} \\ &\ge \frac{nN}{Y_n}(c + 3\delta) - \frac{nN\delta}{Y_n} - \frac{X\|\varphi\|}{Y_n}. \end{aligned}$$

It follows from (3.3) we obtain

$$\lim_{n\to\infty}\frac{nN}{Y_n}=1,\ \lim_{n\to\infty}\frac{X}{Y_n}=0.$$

Thus for sufficiently large n, we have

$$\frac{1}{Y_n}S_{Y_n}\varphi(z)>c+\frac{3}{2}\delta. \tag{3.5}$$

Hence the desired result follows. □

Lemma 3.6. *For sufficiently large n,*

$$\frac{1}{Y_n}\log m\left\{x\in M:\frac{1}{Y_n}S_{Y_n}\varphi(x)>c\right\}\geq h_\nu(f)-\int\psi d\nu-10\gamma-\gamma h_{top}(f).$$

Proof. By lemma 3.5, we have

$$\begin{aligned}&\frac{1}{Y_n}\log m\left\{x\in M:\frac{1}{Y_n}S_{Y_n}\varphi(x)>c\right\}\\ \geq&\frac{1}{Y_n}\log\sum_{z\in F_n}mB_{Y_n}(z,\epsilon')\geq\frac{1}{Y_n}\log\sum_{z\in F_n}C\exp(-S_{Y_n}\psi(z))\\ =&\frac{1}{Y_n}\log\sum_{z\in F_n}\exp(-S_{Y_n}\psi(z))+\frac{1}{Y_n}\log C.\end{aligned}$$

Applying (3.1) and (3.2), we have

$$\begin{aligned}S_{Y_n}\psi(z)<&\sum_{i=1}^{k}\sum_{j=1}^{nNC_i}S_{n_i}\psi(x(i,j))+nN\gamma+X\|\psi\|\\ \leq&\sum_{i=1}^{k}nNC_in_i(\int\psi d\mu_i+\gamma)+nN\gamma+X\|\psi\|\\ =&nN\int\psi d\lambda+2nN\gamma+X\|\psi\|.\end{aligned}$$

It follows from (3.4) lemma 3.3 that

$$\begin{aligned}&\frac{1}{Y_n}\log m\left\{x\in M:\frac{1}{Y_n}S_{Y_n}\varphi(x)>c\right\}\\ \geq&\frac{1}{Y_n}\log\sum_{z\in F_n}\exp\left\{-nN\int\psi d\lambda-2nN\gamma-X\|\psi\|\right\}+\frac{1}{Y_n}\log C\\ \geq&\frac{1}{Y_n}\log\exp\left\{nN(1-\gamma)(h_\lambda(f)-5\gamma)-nN\int\psi d\lambda-2nN\gamma-X\|\psi\|\right\}+\frac{1}{Y_n}\log C\\ =&\frac{1}{Y_n}\left\{nN(h_\lambda(f)-\int\psi d\lambda)+nN(5\gamma^2-7\gamma-\gamma h_\lambda(f))-X\|\psi\|\right\}+\frac{1}{Y_n}\log C\\ \geq&\frac{1}{Y_n}\left\{nN(h_\nu(f)-\int\psi d\nu)+nN(-9\gamma-\gamma h_\lambda(f))-X\|\psi\|\right\}+\frac{1}{Y_n}\log C.\end{aligned}$$

Together with

$$\lim_{n\to\infty}\frac{nN}{Y_n}=1,\ \lim_{n\to\infty}\frac{X}{Y_n}=0,$$

for sufficiently large n, we have

$$\frac{1}{Y_n}\log m\left\{x\in M:\frac{1}{Y_n}S_{Y_n}\varphi(x)>c\right\}\geq h_\nu(f)-\int\psi d\nu-10\gamma-\gamma h_{top}(f).$$

□

For any $n\in\mathbb{N}$, let i_n be the unique natural number such that

$$Y_{i_n}\leq n<Y_{i_n+1}.$$

By (3.5), for sufficiently large n we have

$$\frac{1}{n}S_n\varphi(z)>c+\delta \text{ and } B_n(z,\epsilon')\subseteq\left\{x\in M:\frac{1}{n}S_n\varphi(x)>c\right\}$$

for all $z\in F_{i_n}$. Combining this with previous lemma we obtain

$$\begin{aligned}&\frac{1}{n}\log m\left\{x\in M:\frac{1}{n}S_n\varphi(x)>c\right\}\\ \geq&\frac{Y_{i_n}}{n}\cdot\frac{1}{Y_{i_n}}\log\sum_{z\in F_{i_n}}mB(z,\epsilon')\\ \geq&\frac{Y_{i_n}}{n}\left\{h_\nu(f)-\int\psi d\nu-10\gamma-\gamma h_{top}(f)\right\}.\end{aligned}$$

for sufficiently large n. Since γ was arbitrary, the proof of theorem 2.1 is completed.

4 Some Applications

Example 1 Diffeomorphisms on surfaces Let $f:M\to M$ be a $C^{1+\alpha}$ diffeomorphism with $\dim M=2$ and $h_{top}(f)>0$, then there exists a hyperbolic measure $m\in\mathscr{M}_{\text{erg}}(M,f)$ with Lyapunov exponents $\lambda_1>0>\lambda_2$ (see [19]). If $\beta_1=|\lambda_2|$ and $\beta_2=\lambda_1$, then for any $\epsilon>0$ such that $\beta_1,\beta_2>\epsilon$, we have $m(\Lambda(\beta_1,\beta_2,\epsilon))=1$. Let

$$\widetilde{\Lambda}=\bigcup_{k=1}^{\infty}\text{supp}(m|\Lambda(\beta_1,\beta_2,\epsilon)),$$

then for every $\varphi\in C^0(M)$ and $c\in\mathbb{R}$, the following hold:

(1) For $\psi\in\mathcal{V}^+$, we have

$$\overline{R}(\varphi,[c,\infty))\leq\sup\left\{h_\nu(f)-\int\psi d\nu:\nu\in\mathscr{M}_{\text{inv}}(M,f),\int\varphi d\nu\geq c\right\}.$$

(2) For $\psi \in \mathcal{V}^-$, we have

$$\underline{R}(\varphi,(c,\infty)) \geq \sup\left\{h_\nu(f) - \int \psi d\nu : \mu \in \mathscr{M}_{\text{inv}}(\widetilde{\Lambda}, f), \int \varphi d\nu > c\right\}.$$

Example 2 Nonuniformly hyperbolic systems In [10], Katok described a construction of a diffeomorphism on the 2-torus $\mathbb{T}^2$ with nonzero Lyapunov exponents, which is not an Anosov map. Let f_0 be a linear automorphism given by the matrix

$$A = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$$

with eigenvalues $\lambda^{-1} < 1 < \lambda$. f_0 has a maximal measure μ_1. Let D_r denote the disk of radius r centered at (0,0), where $r > 0$ is small, and put coordinates (s_1, s_2) on D_r corresponding to the eigendirections of A, i.e, $A(s_1, s_2) = (\lambda s_1, \lambda^{-1} s_2)$. The map A is the time-1 map of the local flow in D_r generated by the following system of differential equations:

$$\frac{ds_1}{dt} = s_1 \log\lambda, \frac{ds_2}{dt} = -s_2 \log\lambda.$$

The Katok map is obtained from A by slowing down these equations near the origin. It depends upon a real-valued function ψ, which is defined on the unit interval $[0, 1]$ and has the following properties:

(1) ψ is C^∞ except at 0;

(2) $\psi(0) = 0$ and $\psi(u) = 1$ for $u \geq r_0$ where $0 < r_0 < 1$;

(3) $\psi'(u) > 0$ for every $0 < u < r_0$;

(4) $\int_0^1 \frac{du}{\psi(u)} < \infty$.

Fix sufficiently small numbers $r_0 < r_1$ and consider the time-1 map g generated by the following system of differential equations in D_{r_1}:

$$\frac{ds_1}{dt} = s_1\psi(s_1^2 + s_2^2)\log\lambda, \frac{ds_2}{dt} = -s_2\psi(s_1^2 + s_2^2)\log\lambda.$$

The map f, given as $f(x) = g(x)$ if $x \in D_{r_1}$ and $f(x) = A(x)$ otherwise, defines a homeomorphism of torus, which is a C^∞ diffeomorphism everywhere except for the origin. To provide the differentiability of map f, the function ψ must satisfy some extra conditions. Namely, the integral $\int_0^1 du/\psi$ must converge "very slowly" near the origin. We refer the smoothness to [10]. Here f is contained in the C^0 closure of Anosov diffeomorphisms and even more there is a homeomorphism $\pi : \mathbb{T}^2 \to \mathbb{T}^2$ such that $\pi \circ f_0 = f \circ \pi$. Let $\nu_0 = \pi_* \mu_1$.

In [14], the authors proved that there exist $0 < \epsilon \ll \beta$ and a neighborhood U of ν_0 in $\mathscr{M}_{\rm inv}(\mathbb{T}^2, f)$ such that for any ergodic $\nu \in U$ it holds that $\nu \in \mathscr{M}_{\rm inv}(\widetilde{\Lambda}(\beta,\beta,\epsilon), f)$, where $\widetilde{\Lambda}(\beta,\beta,\epsilon) = \bigcup_{k\geq 1} \mathrm{supp}(\nu_0|\Lambda_k(\beta,\beta,\epsilon))$.

Corollary 4.1. *For every $\varphi \in C^0(\mathbb{T}^2)$ and $c \in \mathbb{R}$, the following hold:*

(1) For $\psi \in \mathcal{V}^+$, we have

$$\overline{R}(\varphi, [c,\infty)) \leq \sup\left\{h_\nu(f) - \int \psi d\nu : \nu \in \mathscr{M}_{\rm inv}(\mathbb{T}^2, f), \int \varphi d\nu \geq c\right\}.$$

(2) For $\psi \in \mathcal{V}^-$, we have

$$\underline{R}(\varphi, (c,\infty)) \geq \sup\left\{h_\nu(f) - \int \psi d\nu : \mu \in \mathscr{M}_{\rm inv}(\widetilde{\Lambda}(\beta,\beta,\epsilon), f), \int \varphi d\nu > c\right\}.$$

In [14], the authors also studied the structure of Pesin set $\widetilde{\Lambda}$ for the robustly transitive partially hyperbolic diffeomorphisms described by Mañé and the robustly transitive non-partially hyperbolic diffeomorphisms described by Bonatti-Viana. They showed that for the diffeomorphisms derived from Anosov systems $\mathscr{M}_{\rm inv}(\widetilde{\Lambda}, f)$ enjoys many members. So our result is applicable to these maps.

Acknowledgements. The research was supported by the National Basic Research Program of China (Grant No. 2013CB834100) and the National Natural Science Foundation of China (Grant No. 11271191).

References

[1] V. Araujo & M.J. Pacifico, Large deviations for non-uniformly expanding maps, *J. Stat. Phys.* **125** (2006), 415-457.

[2] L. Barreira & Y. Pesin, Lyapunov exponents and smooth ergodic theory, Univ. Lect. Series, 23, Providence, RI: American Mathematical Society, (2002).

[3] L. Barreira & Y. Pesin, Nonuniform hyperbolicity: Dynamics of systems with nonzero Lyapunov exponents, Cambridge Press, (2007).

[4] J. Bochi, Genericity of zero Lyapunov exponents, *Ergodic Theory Dynam. Systems* **22** (2002), 1667-1696.

[5] Y.M. Chung, Large deviations on Markov towers, *Nonlinearity* **24** (2011), 1229-1252.

[6] H. Comman & J. Rivera-Letelier, Large deviations principles for non-uniformly hyperbolic rational maps, *Ann. Inst. H. Poincaré. Anal. Non Linéaire* **15** (1998), 539-579.

[7] A. Eizenberg, Y. Kifer & B. Weiss, Large deviations for $\mathbb{Z}^d$-actions, *Commun. Math. Phys.* **164** (1994), 433-454.

[8] R.S. Ellis, Entropy, Large deviations and Statistical Mechanics, Grundlehren der Mathematischen Wissenchaften 271, Spinger-Verlag, New York, (1985).

[9] M. Hirayama, Periodic probability measures are dense in the set of invariant measures, *Discrete Contin. Dyn. Syst.* **9** (2003), 1185-1192.

[10] A. Katok, Bernoulli Diffeomorphisms on Surfaces, *Annals of Math.* (2) **110** (1979), 529-547.

[11] A. Katok, Lyapunov exponents, entropy and periodic points for diffeomorphism, *Publ. Math. IHÉS* **51** (1980), 137-173.

[12] A. Katok & B. Hasselblatt, Introduction to the modern theory of dynamical systems, Encyclopedia of Mathematics and its Applications 54, Cambridge Univ. Press, Cambridge, (1995).

[13] Y. Kifer, Large deviations in dynamical systems and stochastic processes. *Trans. Amer. Math. Soc.* **321(2)** (1990), 505-524.

[14] C. Liang, G. Liao, W. Sun & X. Tian. Saturated sets for nonuniformly hyperbolic systems, arXiv:1110.6091v1.

[15] I. Melbourne & M. Nicol, Large deviations for nonuniformly hyperbolic systems, *Tran. Amer. Math. Soc.* **360** (2008), 6661-6676.

[16] V.I. Oseledec, A multiplicative ergodic theorem, *Trans. Mosc. Math. Soc.* **19** (1969), 197-231.

[17] Y. Pesin, Dimension Theory in Dynamical Systems, Contemporary Views and Applications. Univ. of Chicago Press, (1997).

[18] C.-E. Pfister & W.G. Sullivan, Large deviations estimates for dynamical systems without the specification property. Applications to the β-shifts, *Nonlinearity* **18** (2005), 237-261.

[19] M. Pollicott, Lectures on ergodic theory and Pesin theory on compact manifolds, Cambridge Univ. Press, Cambridge, (1993).

[20] M. Pollicott & R. Sharp, Large deviations for intermittent maps, *Nonlinearity* **22** (2009), 2079-2092.

[21] L. Rey-Bellet & L.-S. Young, Large deviations in non-uniformly hyperbolic dynamical systems, *Ergodic Theory Dynam. Systems* **28(2)** (2008), 587-612.

[22] P. Varandas, Non-uniform specification and large deviations for weak Gibbs measures, *J. Stat. Phys.* **146** (2012), 330-358.

[23] K. Yamamoto, On the weaker forms of specification property and their applications, *Proc. Amer. Math. Soc.* **137** (2009), 3807-3814.

[24] L.-S. Young, Large deviations in dynamical systems, *Trans. Amer. Math. Soc.* **318** (1990), 525-543.

[25] M, Yuri, Large deviations for countable to one Markov systems, *Commun. Math. Phys.* **258(2)** (2005), 455-474.